

## PRODUCT INFORMATION

# BglI

**#ER0071**      2000 u

**Lot:**                      **Expiry Date:**

5'...**G C C N N N N↓N G G C**...3'  
3'...**C G G N↑N N N N C C G**...5'

Concentration:      10 u/μl  
Source:                *Bacillus globigii*  
Supplied with:      1 ml of 10X Buffer O  
                              1 ml of 10X Buffer Tango

**Store at -20°C**



In total 3 vials.

BSA included

[www.thermoscientific.com/fermentas](http://www.thermoscientific.com/fermentas)

## RECOMMENDATIONS

**1X Buffer O** (for 100% BglI digestion)

50 mM Tris-HCl (pH 7.5), 10 mM MgCl<sub>2</sub>, 100 mM NaCl,  
0.1 mg/ml BSA.

**Incubation temperature**

37°C.

**Unit Definition**

One unit is defined as the amount of BglI required to digest 1 μg of lambda DNA in 1 hour at 37°C in 50 μl of recommended reaction buffer.

**Dilution**

Dilute with Dilution Buffer (#B19): 10 mM Tris-HCl (pH 7.4 at 25°C) 100 mM KCl, 1 mM EDTA, 1 mM DTT, 0.2 mg/ml BSA and 50% glycerol.

**Double Digests**

Thermo Scientific Tango Buffer is provided to simplify buffer selection for double digests. 98% of Thermo Scientific restriction enzymes are active in a 1X or 2X concentration of Tango™ Buffer. Please refer to to [www.fermentas.com/doubledigest](http://www.fermentas.com/doubledigest) to choose the best buffer for your experiments.

1X Tango Buffer: 33 mM Tris-acetate (pH 7.9 at 37°C), 10 mM magnesium acetate, 66 mM potassium acetate, 0.1 mg/ml BSA.

## Storage Buffer

BglI is supplied in: 10 mM Tris-HCl (pH 7.5 at 25°C), 300 mM NaCl, 1 mM DTT, 0.1 mM EDTA, 0.2 mg/ml BSA and 50% glycerol.

## Recommended Protocol for Digestion

- Add:

nuclease-free water	16 µl
10X Buffer O	2 µl
DNA (0.5-1 µg/µl)	1 µl
BglI	0.5-2 µl
- Mix gently and spin down for a few seconds.
- Incubate at 37°C for 1-2 hours.

The digestion reaction may be scaled either up or down.

## Recommended Protocol for Digestion of PCR Products Directly after Amplification

- Add:

PCR reaction mixture	10 µl	(~0.1-0.5 µg of DNA)
nuclease-free water	18 µl	
10X Buffer O	2 µl	
BglI	1-2 µl	
- Mix gently and spin down for a few seconds.
- Incubate at 37°C for 1-16 hours.

## Thermal Inactivation

BglI is inactivated by incubation at 65°C for 20 min.

## ENZYME PROPERTIES

### Enzyme Activity in Thermo Scientific REase Buffers, %

B	G	O	R	Tango	2X Tango
0-20	50-100	100	100	0-20	100

## Methylation Effects on Digestion

Dam: never overlaps – no effect.  
Dcm: may overlap – no effect.  
CpG: may overlap – cleavage impaired.  
EcoKI: never overlaps – no effect.  
EcoBI: never overlaps – no effect.

## Stability during Prolonged Incubation

A minimum of 0.1 units of the enzyme is required for complete digestion of 1 µg of lambda DNA in 16 hours at 37°C.

## Digestion of Agarose-embedded DNA

A minimum of 5 units of the enzyme is required for complete digestion of 1 µg of agarose-embedded lambda DNA in 16 hours.

## Number of Recognition Sites in DNA

λ	ΦX174	pBR322	pUC57	pUC18/19	pTZ19R/U	M13mp18/19
29	0	3	2	2	2	1

For **CERTIFICATE OF ANALYSIS** see back page

# CERTIFICATE OF ANALYSIS

## Overdigestion Assay

No detectable change in the specific fragmentation pattern is observed after a 160-fold overdigestion with BglI (10 u/μg lambda DNA x 16 hours).

## Ligation/Recutting Assay

After a 50-fold overdigestion (3 u/μg DNA x 17 hours) with BglI, more than 95% of the digested DNA fragments can be ligated at a 5'-termini concentration of 0.3 μM. More than 95% of these sites can be recut.

## Labeled Oligonucleotide (LO) Assay

No detectable degradation of single-stranded or double-stranded labeled oligonucleotides occurred during incubation with 10 units of BglI for 4 hours.

Quality authorized by:



Jurgita Zilinskiene

## **PRODUCT USE LIMITATION**

This product is developed, designed and sold exclusively *for research purposes and in vitro use only*. The product was not tested for use in diagnostics or for drug development, nor is it suitable for administration to humans or animals.

Please refer to [www.thermoscientific.com/fermentas](http://www.thermoscientific.com/fermentas) for Material Safety Data Sheet of the product.

© 2011 Thermo Fisher Scientific Inc. All rights reserved. All trademarks are the property of Thermo Fisher Scientific Inc. and its subsidiaries.

