

## PRODUCT INFORMATION

# NdeI

**#ER0581**      500 U

**Lot:** \_\_\_\_\_      **Expiry Date:** \_\_\_\_\_

5'...C A↓T A T G...3'

3'...G T A T↑A C...5'

Concentration:      10 U/μL  
Source:              *Neisseria denitrificans*  
Supplied with:      1 mL of 10X Buffer O  
                             1 mL of 10X Buffer Tango

**Store at -20°C**



In total 3 vials.

BSA included

[www.thermoscientific.com/onebio](http://www.thermoscientific.com/onebio)

## RECOMMENDATIONS

**1X Buffer O** (for 100% NdeI digestion)

50 mM Tris-HCl (pH 7.5), 10 mM MgCl<sub>2</sub>, 100 mM NaCl,  
0.1 mg/mL BSA.

**Incubation temperature**

37°C.

**Unit Definition**

One unit is defined as the amount of NdeI required to digest 1 μg of lambda DNA in 1 hour at 37°C in 50 μL of recommended reaction buffer.

**Dilution**

Dilute with Dilution Buffer (#B19): 10 mM Tris-HCl (pH 7.4 at 25°C), 100 mM KCl, 1 mM EDTA, 1 mM DTT, 0.2 mg/mL BSA and 50% glycerol.

**Double Digests**

Thermo Scientific Tango Buffer is provided to simplify buffer selection for double digests. 98% of Thermo Scientific restriction enzymes are active in a 1X or 2X concentration of Tango™ Buffer. Please refer to [www.thermoscientific.com/doubledigest](http://www.thermoscientific.com/doubledigest) to choose the best buffer for your experiments.

1X Tango Buffer: 33 mM Tris-acetate (pH 7.9 at 37°C), 10 mM magnesium acetate, 66 mM potassium acetate, 0.1 mg/mL BSA.

Rev.10

## Storage Buffer

NdeI is supplied in: 10 mM Tris-HCl (pH 7.4 at 25°C), 100 mM KCl, 1 mM DTT, 1 mM EDTA, 0.2 mg/mL BSA and 50% glycerol.

## Recommended Protocol for Digestion

- Add:

|                              |               |
|------------------------------|---------------|
| nuclease-free water          | 16 $\mu$ L    |
| 10X Buffer O                 | 2 $\mu$ L     |
| DNA (0.5-1 $\mu$ g/ $\mu$ L) | 1 $\mu$ L     |
| NdeI                         | 0.5-2 $\mu$ L |
- Mix gently and spin down for a few seconds.
- Incubate at 37°C for 1-16 hours.

The digestion reaction may be scaled either up or down.

## Recommended Protocol for Digestion of PCR Products Directly after Amplification

- Add:

|                      |                                      |
|----------------------|--------------------------------------|
| PCR reaction mixture | 10 $\mu$ L (~0.1-0.5 $\mu$ g of DNA) |
| nuclease-free water  | 18 $\mu$ L                           |
| 10X Buffer O         | 2 $\mu$ L                            |
| NdeI                 | 1-2 $\mu$ L                          |
- Mix gently and spin down for a few seconds.
- Incubate at 37°C for 1-16 hours.

## Thermal Inactivation

NdeI is inactivated by incubation at 65°C for 20 min.

## ENZYME PROPERTIES

### Enzyme Activity in Thermo Scientific REase Buffers, %

| B    | G    | O   | R      | Tango | 2X Tango |
|------|------|-----|--------|-------|----------|
| 0-20 | 0-20 | 100 | 50-100 | 0-20  | 50-100   |

## Methylation Effects on Digestion

Dam: never overlaps – no effect.  
Dcm: never overlaps – no effect.  
CpG: never overlaps – no effect.  
EcoKI: never overlaps – no effect.  
EcoBI: may overlap – no effect.

## Stability during Prolonged Incubation

A minimum of 0.2 units of the enzyme is required for complete digestion of 1  $\mu$ g of lambda DNA in 16 hours at 37°C.

## Digestion of Agarose-embedded DNA

A minimum of 5 units of the enzyme is required for complete digestion of 1  $\mu$ g of agarose-embedded lambda DNA in 16 hours.

## Compatible Ends

Csp6I, FspBI, Tru1I, VspI

## Number of Recognition Sites in DNA

| $\lambda$ | $\Phi$ X174 | pBR322 | pUC57 | pUC18/19 | pTZ19R/U | M13mp18/19 |
|-----------|-------------|--------|-------|----------|----------|------------|
| 7         | 0           | 1      | 1     | 1        | 0        | 3          |

# CERTIFICATE OF ANALYSIS

## Overdigestion Assay

No detectable change in the specific fragmentation pattern is observed after a 160-fold overdigestion with NdeI (10 U/ $\mu$ g lambda DNA  $\times$  16 hours).

## Ligation and Recleavage (L/R) Assay

The ligation and recleavage assay was replaced with LO test after validating experiments showed LO test ability to trace nuclease and phosphatase activities with sensitivity that is higher than L/R by a factor of 100.

## Labeled Oligonucleotide (LO) Assay

No detectable degradation of single-stranded or double-stranded labeled oligonucleotides occurred during incubation with 10 units of NdeI for 4 hours.

Quality authorized by:



Jurgita Zilinskiene

## **PRODUCT USE LIMITATION**

This product is developed, designed and sold exclusively *for research purposes and in vitro use only*. The product was not tested for use in diagnostics or for drug development, nor is it suitable for administration to humans or animals.

Please refer to [www.thermoscientific.com/onebio](http://www.thermoscientific.com/onebio) for Material Safety Data Sheet of the product.

© 2012 Thermo Fisher Scientific Inc. All rights reserved. All trademarks are the property of Thermo Fisher Scientific Inc. and its subsidiaries.